

ORIGINAL ARTICLE

Distribution and variability of deformed wing virus of honeybees (*Apis mellifera*) in the Middle East and North Africa

Nizar Jamal Haddad¹, Adjlane Nouredine², Banan Al-Shagour¹, Wahida Loucif-Ayad³, Mogbel A. A. El-Niweiri⁴, Eman Anaswah¹, Wafaa Abu Hammour¹, Dany El-Obeid⁵, Albaba Imad⁶, Mohamed A. Shebl⁷, Abdulhusien Sehen Almaleky⁸, Abdullah Nasher⁹, Nagara Walid¹⁰, Mohamed Fouad Bergigui¹¹, Orlando Yañez¹² and Joachim R. de Miranda¹³

¹Bee Research Department, National Center for Agriculture Research and Extension, Baq'a, Jordan; ²Department of Biology, M'hamed Bougara University of Boumerdes, ENS Kouba, Algeria; ³Laboratory of Applied Animal Biology, University Badji-Mokhtar, Annaba, Algeria; ⁴Department of Bee Research, Environment, Natural Resources and Desertification Research Institute, National Centre for Research, Khartoum, Sudan; ⁵Faculty of Agriculture and Veterinary Sciences, Lebanese University, Beirut, Lebanon; ⁶West Bank, State of Palestine, Halhul-Hebron District, Palestine; ⁷Department of Plant Protection, Suez Canal University, Ismailia, Egypt; ⁸Extension Department, Qadysia Governate Agricultural Directorate, Iraq; ⁹Department of Plant Protection, Sana'a University, Sana'a, Yemen; ¹⁰National Federation of Tunisian beekeepers, Tunis, Tunisia; ¹¹Ruchers El Bakri, Hay Assalam-Sidi Slimane, Rabat, Morocco; ¹²Institute of Bee Health, Vetsuisse Faculty, University of Bern, Bern, Switzerland and ¹³Department of Ecology, Swedish University of Agricultural Sciences, Uppsala, Sweden

Abstract Three hundred and eleven honeybee samples from 12 countries in the Middle East and North Africa (MENA) (Jordan, Lebanon, Syria, Iraq, Egypt, Libya, Tunisia, Algeria, Morocco, Yemen, Palestine, and Sudan) were analyzed for the presence of deformed wing virus (DWV). The prevalence of DWV throughout the MENA region was pervasive, but variable. The highest prevalence was found in Lebanon and Syria, with prevalence dropping in Palestine, Jordan, and Egypt before increasing slightly moving westwards to Algeria and Morocco. Phylogenetic analysis of a 194 nucleotide section of the DWV Lp gene did not identify any significant phylogenetic resolution among the samples, although the sequences did show consistent regional clustering, including an interesting geographic gradient from Morocco through North Africa to Jordan and Syria. The sequences revealed several clear variability hotspots in the deduced amino acid sequence, which furthermore showed some patterns of regional identity. Furthermore, the sequence variants from the Middle East and North Africa appear more numerous and diverse than those from Europe.

Key words *Apis mellifera*; deformed wing virus; Middle East; North Africa; prevalence; variability

Introduction

Honeybees (*Apis mellifera* L.) play an important role in nature and agriculture, where they are needed for the

pollination of numerous crops. In recent years, there have been increasing reports of substantial losses of honeybee colonies in many countries (vanEngelsdorp & Meixner, 2010). These mortalities have multiple causes, from the still mysterious rapid adult depopulation phenomena (vanEngelsdorp *et al.*, 2009, 2011) and acute poisoning by novel insecticides (Rosenkranz *et al.*, 2010; Blacquiere *et al.*, 2012; Lu *et al.*, 2014) to the more common mortalities due to honeybee parasites and pathogens, such

Correspondence: Nizar Jamal Haddad, Bee Research Department, National Center for Agriculture Research and Extension, P.O. Box 639, Baq'a 19381, Jordan. Tel: +96 2777327437; fax: +96 265372516; email: drnizarh@gmail.com

as Nosema (Fries, 2010), foulbrood (Forsgren & Fries, 2010) and in particular the parasitic mite *Varroa destructor* (Rosenkranz et al., 2010) and the lethal virus epidemics it initiates and vectors (de Miranda & Genersch, 2010; de Miranda et al., 2010).

In the Arab world, abnormal mortality and local disappearances of bees are often reported by beekeepers (Haddad, 2011; Adjlane et al., 2012a,b) and there has been considerable recent research in the region on the prevalence of *Varroa* (El-Niweiri & El-Sarrag, 2006; Loucif Ayad et al., 2008; Adjlane, 2010; Belaid & Et Doumandji, 2010), Nosema, American foulbrood (AFB), the tracheal mite *Acarapis woodi* (El-Niweiri et al., 2008; Adjlane et al., 2012a,b; Haddad, 2014) and honeybee viruses (Haddad et al., 2008; Loucif-Ayad et al., 2013; Adjlane & Haddad, 2013) to complement historical records (summarized by Ellis & Munn, 2005). However, there have been very few studies on viral diseases of honeybees that cover the entire Middle East and North Africa region (hereafter referred to as MENA region) and none concerning the genetic character and diversity of the viruses in such colonies.

The principal virus associated with elevated colony mortality is *deformed wing virus* (DWV). This virus was first isolated in the 1970's from a sample of asymptomatic dead honeybees from Egypt, and called Egypt bee virus (EBV), before it was found to be serologically related to a virus from DWV-symptomatic bees from Japan, isolated in the early 1980's (Bailey & Ball, 1991).

DWV has become one of the major virus threats to the honeybee industry due to its synergism with *V. destructor* (Martin, 2001), which acts as an inducer and highly efficient vector of the virus (Bowen-Walker et al., 1999; Nordström, 2000; de Miranda & Genersch, 2010). The mites cause immunosuppression in the bees (Yang & Cox-Foster, 2005), activating inapparent DWV infections which are subsequently transmitted extremely efficiently by the mite, precipitating a lethal epidemic that increasingly also involves other transmission routes (de Miranda & Genersch, 2010). Even when it is not a direct vector, *Varroa* can increase the susceptibility of honeybees to other opportunistic pathogens (Yang & Cox-Foster, 2005), leading to a complex disease profile at colony level (Hung & Shimanuki, 1999; Tentcheva et al., 2004a,b; de Miranda et al., 2012), and a progressive reduction in colony performance, followed by large-scale death. In the absence of mite, DWV is largely innocuous and often even undetectable, such that effective mite control stops and even reverses the DWV epidemic (Martin et al., 2010; Rosenkranz et al., 2010).

Typical symptoms exhibited by severely DWV-infected bees are crumpled wings, bloated abdomens, paralysis, learning impairment, and a drastically shortened life span

(Bailey & Ball, 1991). DWV has been detected in all honeybee life stages, as well as glandular secretions (Chen et al., 2005; Yue & Genersch, 2005), through which it is orally transmitted to larvae. DWV is also infectious in bumble bees, where it causes similar symptoms (Genersch et al., 2006), and it is an apiculture-derived threat to wild pollinators (Singh et al., 2010; Fürst et al., 2014).

The genomes of DWV and its close relative *Varroa destructor virus-1* (VDV-1) were published in the mid 2000's (Fujiyuki et al., 2004; Ongus et al., 2004; Lanzi et al., 2006). The genome is 10 140 nucleotides long, excluding the poly-A tail, and contains a single large open reading frame encoding a 328-kilo Dalton (kDa) polyprotein. The coding region is flanked on the 5' and 3' sides by untranslated regions (UTR) of about 1140 and 317 nucleotides, containing sequences and RNA secondary structures for replication and translation (Lanzi et al., 2006; de Miranda & Genersch, 2010). Like all viruses, DWV exists as a dynamic population of variants, recombinants and mutations that provide the raw material for rapid adaptation, and that is constantly replenished through high rates of mutation and recombination (de Miranda & Genersch, 2010; Moore et al., 2011; Domingo, 2012; Zioni et al., 2012; Wang et al., 2013; Ryabov et al., 2014). Individual DWV variants have been linked to elevated aggression in bees (Fujiyuki et al., 2004) although DWV is of course not the only, or even the main factor in aggressive behavior (Rortais et al., 2006; Terio et al., 2008). Other individual variants or recombinants have been linked to molecular virulence, that is, high virus titres (Ryabov et al., 2014), and at bee population level there is evidence for a constriction of DWV genetic variation and the persistence of certain variants during successive mite invasions (Martin et al., 2012), suggesting a degree of adaptation to *Varroa*-mediated transmission and epidemiology. In addition to the specific traits of individual strains, virus variability itself can influence virulence and associated parameters, with more variable isolates often also more virulent (Lauring & Andino, 2010).

Despite the severity and impact of DWV in the MENA region, little is known about its genetic diversity and history in the region. This is the first study of the prevalence, phylogenetic relatedness and variability of DWV isolates collected from the MENA region.

Materials and methods

Sample collection

Three hundred and eleven adult worker bee samples were collected during 2012 from 143 apiaries located in 12 countries, distributed as follows: Lebanon (9), Syria




Fig. 1 Origin of samples and deformed wing virus (DWV) prevalence in the MENA region countries. Red dots denote positive samples, MENA apiaries (note that the map has insufficient resolution for uniquely marking all apiaries); blue dots mark the origin of European DWV sequences used in the phylogenetic and variability analysis. The number of samples from each country is indicated within the respective prevalence pie-chart. The number of apiaries per country is presented in the table within the figure.

(19), Iraq (18), Palestine (10), Jordan (26), Egypt (10), Libya (3), Tunisia (5), Algeria (21), Morocco (9), Yemen (9), and Sudan (4), and with between 1 and 5 colonies sampled per apiary. The approximate locations of the sampled apiaries are shown in Figure 1. The colonies and apiaries sampled were usually *Varroa*-infested and the bee samples included both bees with and without typical pathological symptoms of deformed wing virus infection. Each sample consisted of 10–15 bees, collected from inside the colonies, preserved in >20 mL 100% alcohol (Evans *et al.*, 2013) and shipped at ambient temperature to Jordan to arrive within 3 weeks of collection. Upon arrival, the bees were removed from the alcohol and stored at –80 °C until RNA extraction. Ethanol was chosen as the most suitable preservative for the realities of the sampling conditions in the region.

This sample collection and preservation strategy, although perhaps not optimal from a purely molecular perspective (Chen *et al.*, 2007; Evans *et al.*, 2013), nevertheless represented the best overall strategy available for the modest objectives of the study, the scale of the study, the materials available and the local realities across the entire MENA region. This placed limits on the types of assay that can be reliably conducted on the samples, excluding those requiring highly intact RNA (Bustin *et al.*, 2009; Dainat *et al.*, 2011), as well as on the interpretation of the data.

RNA extraction

Five bees from each sample were homogenized in bulk using standard methods (de Miranda *et al.*, 2013) and total RNA was purified from the homogenate using the RNeasy plant extraction kit (Qiagen, ~~Qiagen~~, Hilden, Germany). The approximate RNA concentration of each sample was determined using the NanoDrop 2000 (Thermo Scientific, ~~Qiagen~~, Hilden, Germany), and the RNA samples were stored at –80 °C until further use.

RT-PCR

A simple, qualitative RT-PCR assay was chosen as the most appropriate DWV diagnostic tool for the sampling strategy employed, and the objectives of the study. The assay was based on a small (194 nucleotide) PCR fragment, to minimize the effects of possible RNA degradation on the ability to reliably detect DWV RNA, and thereby reduce the risk of false-negative data. One-step RT-PCR was performed as described by Genersch (2005) using a one-step RT-PCR kit (Qiagen) following the manufacturer's instructions. Each reaction contained 0.5 ng total RNA, 0.1 mmol/L of forward

(5' CTTACTCTGCCGTCGCCCA 3') and reverse (5' CCGTTAGGAAGCTCATTATCGCG 3') primers in 50 µL total volume. These primers amplify a 194 nucleotide section of the Lp gene (Lanzi *et al.*, 2006), at the start of the main open reading frame, which is well-suited for DWV variability studies (de Miranda & Genersch, 2010). These primers are designed conservatively to allow the detection of most DWV variants in the sequence space between DWV and its closest relative, VDV-1 (Ongus *et al.*, 2004), thereby also reducing the risk of false-negative data, in this case through DWV sequence divergence. The following cycling conditions were used: 30-min cDNA synthesis at 50 °C, 10-min denaturation at 95 °C followed by 40 cycles of: 30 sec at 94 °C, 30 sec at 65 °C, 30 sec at 72 °C, followed by a final elongation step of 7 min at 72 °C. No template controls (NTC) were included in all assays. Ten microliter of each reaction was analyzed on a 1.5% agarose gel containing ethidium bromide and visualized on a UV trans-illuminator to identify the 194 bp DWV-specific RT-PCR product produced by the assay.

Sequencing

Sixty-seven RT-PCR products out of 119 DWV-positive samples, distributed equitably between countries and apiaries, were selected for sequencing: Lebanon (3), Syria (13), Iraq (1), Palestine (0), Jordan (16), Egypt (3), Libya (0), Tunisia (4), Algeria (11), Morocco (11), Yemen (5). The PCR products were gel purified using MEGA quick-spin (~~Kyeonggi-do, Korea~~)TM PCR and Agarose Gel DNA Extraction System (Kyeonggi-do, Korea) according to the protocol recommended by the manufacturer. The purified RT-PCR products were sequenced in both orientations using the primers used for amplification. The forward and reverse sequences of each PCR product were combined into a single sequence. Any conflicts between the forward and reverse sequences, as well as instances of true polymorphism within the PCR product (Forsgren *et al.*, 2009), were resolved by visual examination of the sequence electropherograms. The MENA sequences were submitted to GenBank under accession numbers KT591877–KT591943.

Phylogenetic and variability analyses

The sequences were aligned to homologous sequences found in the public nucleic acid databases using CLUSTAL-omega, at www.ebi.ac.uk (Sievers *et al.*, 2011; McWilliam *et al.*, 2013). This added one sequence from Israel/Palestine (GenBank accession JF440526; Zioni *et al.*, 2011) to the MENA dataset, plus a large set

of Swedish, European and Far East Asian sequences (GenBank accessions JF346550–JF346657; previously described in Forsgren *et al.*, 2009 and de Miranda & Genersch, 2010) for comparison. Phylogenetic analysis was performed in MEGA6 (Tamura *et al.*, 2013) using Minimum Evolution criteria. The robustness of the partitions was tested by bootstrap analysis (500 bootstraps). The variability among the DWV sequences was analyzed by exporting the translated sequences to Excel, tabulating the prevalence of the different amino acid variants at each position in different geographically-defined groups of sequences (34 each for the Middle East and North Africa & Yemen; 31 for Europe, 96 for Sweden, and 21 for a range of Far East Asian sequences) and converting this information to basic stacked histograms.

Results

DWV prevalence

The survey results showed that DWV is present throughout the MENA region, with the possible exception of Sudan (Fig. 1) and that there is considerable variation in DWV prevalence between the MENA countries. A few patterns can be discerned: the highest prevalence was found in Syria and Lebanon, with a lower prevalence in Palestine, Jordan, Iraq, and Yemen. This contrasts with a much lower prevalence in Egypt, Libya, and Tunisia with a gradual increase again when moving westwards toward Algeria and Morocco (Fig. 1). These prevalence estimates are conservative (i.e., minimum) due to the risk of false-negative results, either because of RNA degradation (a risk of the sample preservation strategy) or because the local DWV variants lie outside the specificity of the DWV assay (see de Miranda *et al.*, 2013). Such risk is part of any sampling strategy and was minimized here by designing a simple, small PCR assay with highly conserved primers. The apparent absence of DWV from Sudan in particular should be interpreted with caution, since this result is based on just 4 samples.

Phylogenetic and variability analyses

The phylogenetic analysis of the nucleotide sequences reveals consistent regional clustering (Fig. 2), including an interesting geographic gradient from Morocco through North Africa to Jordan and Syria. Unfortunately; there is very little reliable resolution throughout the phylogeny except for the major branches, making it difficult to draw definitive conclusions from these trends in the data. A different approach to the data is to look at the patterns

of DWV variability between different regions (Fig. 3). There are clear hotspots in the amino acid sequence, where variants are more frequent and more diverse. The same hotspots, and variants, appear in all regions, though in different combinations and frequencies. A particularly variable section of the sequence is between amino acids 38 and 52, where there are both numerous alternative variants at individual positions and different blends of variants in different geographic regions. Particular variants at individual positions can be identified with particular countries within each region (Fig. 3). However, such variants are rarely consensual for a region and it is more the frequency and distribution of variants that is characteristic, rather than the variant itself. The other observation is that the sequence variants from the MENA region appear more numerous and diverse than the variants from Europe.

Discussion

DWV had previously detected in Turkey (Gülmez *et al.*, 2009) and Israel (Zioni *et al.*, 2011) as well as in Egypt, Tunisia, and Saudi Arabia (Ellis & Munn, 2005). While neither *Varroa* nor DWV could be detected in a recent survey of Ugandan bees (Kajobe *et al.*, 2010), both were detected frequently in Kenya (Muli *et al.*, 2014) while *Varroa* has been in Sudan since 2006 at least (El-Niweiri & El-Sarrag, 2006; El-Niweiri *et al.*, 2008) and has also been detected in Ethiopia, together with symptoms typical of DWV infection (Begna, 2015).

The high prevalence of DWV throughout the MENA region is in itself not surprising: it merely confirms the known history of *V. destructor* infestation in the region, which is the primary driver of DWV epidemiology (de Miranda & Genersch, 2010). It is the variation in prevalence between the countries that is perhaps more informative, since these may be related to the *Varroa* invasion history of the different countries: the countries along the Red Sea and in East African are thought to have been invaded relatively recently, with limited impact and possibly positive relationship between geographic elevation and the severity of varroasis (Muli *et al.*, 2014). This contrasts with the much longer history of *Varroa* infestation in the Middle East, where *Varroa* was first detected in the 1980's in Jordan, and caused heavy losses to the commercial and traditional colonies (Ministry of Agriculture, 1986), most likely due to DWV transmission (Haddad *et al.*, 2008). Regional and national differences in *Varroa* management strategies may also have affected the prevalence patterns. Similar observations of the progress of disease and colony losses following the *Varroa* infestation front have been made in other countries (Bailey & Ball, 1991; Shimanuki




Fig. 2 Phylogram of the genetic relationships between the DWV isolates from the MENA region (current study), Europe, and Asia, as determined from a 194 nucleotide fragment of the DWV Lp gene. The statistical strength of the nodes is shown as the percentage of correct partitions in a 500 replicate bootstrap analysis. Only bootstrap values >65% are shown. Different geographic regions are identified by different colored circles.

et al., 1994; Topley *et al.*, 2005; Dainat *et al.*, 2008; Martin *et al.*, 2013), including the recent *Varroa* invasions in Hawaii, USA (Martin *et al.*, 2012) and New Zealand (Stevenson *et al.*, 2005; Mondet *et al.*, 2014).

Likewise, the documentation of various DWV genetic microvariants throughout the MENA region is itself not particularly informative: such differences are expected when surveying broad geographic regions with a large degree of isolation. Rather, it is the trends in the DWV phylogenetic and variability data that are of interest, with

lower DWV variability in the regions with a longer history of *Varroa* infestation (see also Martin *et al.*, 2012), and a phylogenetic clustering pattern that matches this geographic trend, with the more distally clustering variants from regions with longer *Varroa* invasion history (and perhaps longer evolutionary isolation) with the more proximally clustering variants in regions with a more recent invasion history.

It should be stressed here that none of the phylogenetic clustering patterns are statistically significant, and




Fig. 3 Distribution and frequency of the DWV amino acid sequence variants within different geographic regions, based on the 194 nucleotide sequences of the DWV Lp-gene. The consensus amino acid sequence and the variant amino acids at each position are shown below the histogram. Those amino acid variants appearing only once in the entire dataset given in gray type, those appearing more than once are in black type. At each position the primary alternative amino acid variants are colored blue in the histogram, followed by red, green, and purple for the 3rd, 4th, and 5th variants.

that both the tree, the variability data and the prevalence data are equally supportive of a number of alternative *Varroa* and/or DWV invasion scenarios, not necessarily purely linear from the Middle East westwards through North Africa. Importation and regional trade in bees and bee products is an efficient means of spreading disease agents and could easily have contributed to the data patterns found here. Such conclusions, as well as possible links between virus variants/variability and molecular or biological virus virulence, require more precise and dedicated studies.

Conclusion

The deformed wing virus (DWV) is present throughout the Middle East and North Africa (MENA) region, with discernible trends and variation across the region in DWV prevalence, nucleotide sequence and genetic variability, despite the absence of statistical significance to most of the differences between regions and countries. This is the first study of its kind that reports the prevalence of DWV in honeybees across the MENA region countries, and that evaluates the variability of DWV sequences within the

MENA region and compares this to the north Mediterranean basin and Europe.

Acknowledgments

Authors are grateful to collaborators who helped us in sampling: Taoufik Ben Hamida from Tunisia, Prof. Husan Mustafa from Egypt. Special thanks to collaborators who collected samples under very difficult circumstances: Ibrahim Majid, Sabah, Montasir, and Ahmed Najah from Iraq; Mustafa El-Fituri from Libya; Mayyada Haj Ali from Syria. We also appreciate the efforts of Ahmed Batain and Moath Al-Gharaibeh from Jordan for their technical field and laboratory support. The authors would like to thank USAID M32-035 and USAID M29-075 project for funding this project.

Disclosure

The authors declare no conflicts of interest, or specific financial interests, relationships or affiliations.

Authors' contributions

Designed, fundraising, and coordinated the experiments (NJH); provided samples' data (All); conducted the experiments (NJH, BAS, EA); analyzed the data (NJH, JdM, OY, WH); prepared the manuscript (NJH, JdM, OY, AN). All authors read and approved the final manuscript.

References

- Adjlane, N. (2010) Association of oxalic and lactic acid for *Varroa* control in Algeria. *First World Conference on Organic Beekeeping*, 27–29 août 2010, Sunny Beach, p. 52.
- Adjlane, N., Doumandji, S.E. and Haddad, N. (2012a) Situation de l'apiculture en Algérie: facteurs menaçant la survie des colonies d'abeilles locales *Apis mellifera intermissa*. *Cahiers Agricultures*, 21, 235–241.
- Adjlane, N., Kechih, S., Doumandji, S.E. and Haddad, N. (2012b) Survey of American foulbrood in *Apis mellifera intermissa* colonies in mid-northern region of Algeria. *Uludag Bee Journal*, 12, 98–105.
- Adjlane, N. and Haddad, N. (2013) Deformed wing virus and *Varroa destructor* in the local honey bee colonies *Apis mellifera intermissa* in Algeria. World Academy of Science, Engineering and Technology, International Science Index. *Agricultural and Biosystem Engineering*, 1(9), 22.
- Bailey, L. and Ball, B.V. (1991) *Honey Bee Pathology*. London, Academic Press.
- Begna D. (2015) Occurrences and distributions of honeybee (*Apis mellifera Jemenetica*) varroa mite (*Varroa destructor*) in Tigray Region, Ethiopia. *Journal of Fisheries Livestock Production*, 3, 1–4.
- Belaid, M. and Et Doumandji, S. (2010) Effet du *Varroa destructor* sur la morphométrie alaire et sur les composants du système immunitaire de l'abeille ouvrière *Apis mellifera intermissa*. *Lebanese Science Journal*, 11, 45–53.
- Blacquiere, T., Smaghe, G., van Gestel, C.A.M. and Mommaerts, V. (2012) Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology*, 21, 973–992.
- Bowen-Walker, P.L., Martin, S.J. and Gunn, A. (1999) The transmission of deformed wing virus between honeybees (*Apis mellifera* L.) by the ectoparasitic mite *Varroa jacobsoni* Oud. *Journal of Invertebrate Pathology*, 73, 101–106.
- Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L., Vandesompele, J. and Wittwer, C.T. (2009) The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry*, 55, 611–622.
- Chen, Y.P., Higgins, J.A. and Feldlaufer, M.F. (2005) Quantitative real-time reverse transcription-PCR analysis of deformed wing virus infection in the honeybee (*Apis mellifera* L.). *Applied and Environmental Microbiology*, 71, 436–441.
- Chen, Y.P., Evans, J.D., Hamilton, M. and Feldlaufer, M.F. (2007) The influence of RNA integrity on the detection of honey bee viruses: molecular assessment of different sample storage methods. *Journal of Apicultural Research*, 46, 81–87.
- Dainat, B., Imdorf, A., Charrière, J.D. and Neumann, P. (2008) Virus des abeilles: revue des connaissances actuelles. *Revue Suisse d'apiculture*, 129, 8–13.
- Dainat, B., Evans, J.D., Chen, Y.P. and Neumann, P. (2011) Sampling and RNA quality for diagnosis of honey bee viruses using quantitative PCR. *Journal of Virological Methods*, 174, 150–152.
- de Miranda, J.R., Bailey, L., Ball, B.V., Blanchard, P., Budge, G.E., Chejanovsky, N., Chen, Y.P., Gauthier, L., Genersch, E., de Graaf, D.C., Ribière, M., Ryabov, E., De Smet, L. and van der Steen, J.J.M. (2013) Standard methods for virus research in *Apis mellifera*. *The COLOSS BEEBOOK, Volume II: Standard Methods for Apis Mellifera Pest and Pathogen Research* (eds. V. Dietemann, J.D. Ellis & P. Neumann). *Journal of Apicultural Research*, 52, 1–55.
- de Miranda, J.R. and Genersch, E. (2010) Deformed wing virus. *Journal of Invertebrate Pathology*, 103, 48–61.
- de Miranda, J.R., Cordonni, G. and Budge, G. (2010) The acute bee paralysis virus-Kashmir bee virus-Israeli acute paralysis virus complex. *Journal of Invertebrate Pathology*, 103, S30–S47.

- de Miranda, J.R., Gauthier, L., Ribiere, M. and Chen, Y.P. (2012) Honey bee viruses and their effect on bee and colony death. *Honey Bee Colony Health: Challenges and Sustainable Solutions* (eds. D. Sammatro & Y.A. Yoder), pp. 71–102. Taylor & Francis Group Ltd, Oxford, England.
- Domingo, E., Sheldon, J. and Perales, C. (2012) Viral quasispecies evolution. *Microbiology and Molecular Biology Reviews*, 76, 159–216.
- Ellis, J.D. and Munn, P.A. (2005) The worldwide health status of honey bees. *Bee World*, 86, 88–101.
- El-Niweiri, M.A.A. and El-Sarrag, M.S.A. (2006) Detection of the parasitic mite (*Varroa jacobsoni*) of honeybees *Apis mellifera* in Sudan. *Albuhuth*, 10, 60–75.
- El-Niweiri, M.A.A., El-Sarrag, M.S.A. and Satti, A.A. (2008) Survey of diseases and parasites of honey bees (*Apis mellifera* L.) in Sudan. *Sudan Journal of Basic Sciences*, 14, 141–159.
- Evans, J.D., Schwarz, R.S., Chen, Y.P., Budge, G., Cornman, R.S., De La Rua, P., de Miranda, J.R., Foret, S., Foster, L., Gauthier, L., Genersch, E., Gisder, S., Jarosch, A., Kucharski, R., Lopez, D., Lun, C.M., Moritz, R.F.A., Maleszka, R., Muñoz, I. and Pinto, M.A. (2013) Standard methods for virus research in *Apis mellifera*. *The COLOSS BEEBOOK, Volume I: Standard Methods for Apis mellifera Research* (eds. V. Dietemann, J.D. Ellis & P. Neumann). *Journal of Apicultural Research*, 52, 1–3.
- Forsgren, E., de Miranda, J.R., Isaksson, M., Wei, S. and Fries, I. (2009) Deformed wing virus associated with *Tropilaelaps mercedesae* infesting European honey bees (*Apis mellifera*). *Experimental and Applied Acarology*, 47, 87–97.
- Forsgren, E. and Fries, I. (2010) Comparative virulence of *Nosema ceranae* and *Nosema apis* in individual European honey bees. *Veterinary Parasitology*, 170(3–4), 212–217.
- Fries, I. (2010) *Nosema ceranae* in European honey bees. *Journal of Invertebrate Pathology*, 103, S73–S79.
- Fujiyuki, T., Takeuchi, H., Ono, M., Ohka, S., Sasaki, T., Nomoto, A. and Kubo, T. (2004) Novel insect picorna-like virus identified in the brains of aggressive worker honeybees. *Journal of Virology*, 78, 1093–1100.
- Fürst, M.A., McMahon, D.P., Osborne, J.L., Paxton, R.J. and Brown, M.J.F. (2014) Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature*, 506, 364–366.
- Genersch, E. (2005) Development of a rapid and sensitive RT-PCR method for the detection of deformed wing virus: a pathogen of the honey bee (*Apis mellifera*). *Veterinary Journal*, 169, 121–123.
- Genersch, E., Yue, C., Fries, I. and de Miranda, J.R. (2006) Detection of deformed wing virus, a honey bee viral pathogen, in bumble bees (*Bombus terrestris* and *Bombus pascuorum*) with wing deformities. *Journal of Invertebrate Pathology*, 91, 61–63.
- Gülmez, Y., Bursalı, A. and Tekin, S. (2009) First molecular detection and characterization of deformed wing virus (DWV) in honeybees (*Apis mellifera* L.) and mite (*Varroa destructor*) in Turkey. *African Journal of Biotechnology*, 8, 3698–3702.
- Haddad, N. (2011) Bee losses and colony collapse disorder in the Middle East. *Uludag Bee Journal*, 11, 17–24.
- Haddad, N. (2014) First detection of *Nosema ceranae* in Jordan. *European Journal of Scientific Research*, 10(35), 91–96.
- Haddad, N., Brake, M., Megrade, H. and De Miranda, J. (2008) The first detection of honeybee viral diseases in Jordan using the RT-PCR. *Jordan Journal of Agricultural Sciences*, 4, 57–61.
- Hung, A.C.F. and Shimanuki, H. (1999) A scientific note on the detection of Kashmir bee virus in individual honeybees and *Varroa jacobsoni* mites. *Apidologie*, 30, 353–354.
- Kajobe, R., Marris, G., Budge, G., Laurenson, L., Cordoni, G., Jones, Wilkins, S., Cuthbertson, A.G.S. and Brown, A.M. (2010) First molecular detection of a viral pathogen in Ugandan honeybees. *Journal of Invertebrate Pathology*, 104, 153–156.
- Lanzi, G., de Miranda, J.R., Boniotti, M.B., Cameron, C.E., Lavazza, A., Capucci, L., Camazine, S.M. and Rossi, C. (2006) Molecular and biological characterization of deformed wing virus of honeybees (*Apis mellifera* L.). *Journal of Virology*, 80, 4998–5009.
- Lauring, A.S. and Andino, R. (2010) Quasi-species theory and the behavior of RNA viruses. *PLoS Pathogens*, 6, e1001005.
- Lu, C.S., Warchol, K. and Callahan, R. (2014) Sub-lethal exposure to neonicotinoids impaired honey bees winterization before proceeding to colony collapse disorder. *Bulletin of Insectology*, 67, 125–130.
- Loucif-Ayad, W., Aribi, N., Smagghe, G. and Oltani, N. (2008) Evaluation of secondary effects of some Acaricides on *Apis mellifera intermissa* (Hymenoptera, Apidae): acetylcholinesterase and glutathione S-transferase activities. *European Journal of Scientific Research*, 21, 642–649.
- Loucif-Ayad, W., Chefrour, A., Algharibeh, M. and Haddad, N. (2013) First detection of deformed wing virus of honeybees in Algeria. *Phytoparasitica*, 4, 445–447.
- Martin, S.J. (2001) The role of *Varroa* and viral pathogens in the collapse of honeybee colonies: a modelling approach. *Journal of Applied Ecology*, 38, 1082–1093.
- Martin, S.J., Highfield, A.C., Bretell, L., Villalobos, E.M., Budge, G.E., Powell, M., Nikaido, S. and Schroeder, D.C. (2012) Global honey bee viral landscape altered by a parasitic mite. *Science*, 336, 1304–1306.
- Martin, S.J., Ball, B.V. and Carreck, N. (2010) Prevalence and persistence of deformed wing virus (DWV) in untreated or acaricide-treated *Varroa destructor* infested honey bee (*Apis mellifera*) colonies. *Journal of Apicultural Research*, 42, 72–79.

- Martin, S.J., Ball, B.V. and Carreck, N. (2013) The role of deformed wing virus in the initial collapse of varroa infested honey bee colonies in the UK. *Journal of Apicultural Research*, 52, 251–258.
- McWilliam, H., Li, W., Uludag, M., Squizzato, S., Park, Y.M., Buso, N., Cowley, A.P. and Lopez, R. (2013) Analysis tool web services from the EMBL-EBI. *Nucleic Acids Research*, 41, W597–W600.
- Ministry of Agriculture of Jordan (1986) Department of agricultural statistics. *Beekeeping Statistical Book*.
- Mondet, F., de Miranda, J.R., Kretschmar, A., Le Conte, Y. and Mercer, A.R. (2014) On the front line: quantitative virus dynamics in honeybee (*Apis mellifera* L.) colonies along a new expansion front of the parasite *Varroa destructor*. *PLoS Pathogens*, 10, e1004323.
- Moore, J., Jironkin, A., Chandler, D., Burroughs, N. and Evans, D.J. (2011) Recombinants between deformed wing virus and *Varroa destructor* virus-1 may prevail in *Varroa destructor*-infested honeybee colonies. *Journal of General Virology*, 92, 156–161.
- Muli, E., Patch, H., Frazier, M., Frazier, J., Torto, B., Baumgarten, T., Kilonzo, J., Kimani, J.N., Mumoki, F., Masiga, D., Tumlinson, J. and Grozinger, C. (2014) Evaluation of the distribution and impacts of parasites, pathogens, and pesticides on honey bee (*Apis mellifera*) populations in East Africa. *PLoS ONE*, 9, e94459.
- Nordström, S., 2000. Virus infections and varroa mite infestations in honey bee colonies. *PhD thesis*. Swedish University of Agricultural Sciences, Uppsala, Sweden.
- Ongus, J.R., Peters, D., Bonmatin, J.M., Bengsch, E., Vlak, J.M. and van Oers, M.M. (2004) Complete sequence of a picorna-like virus of the genus *Iflavirus* replicating in the mite *Varroa destructor*. *Journal of General Virology*, 85, 3747–3755.
- Rortais, A., Tentcheva, D., Papachristoforou, A., Gauthier, L., Arnold, G., Colin, M.E. and Bergoin, M. (2006) Deformed wing virus is not related to honey bees aggressiveness. *Virology Journal*, 3, e61.
- Rosenkranz, P., Aumeier, P. and Ziegelmann, B. (2010) Biology and control of *Varroa destructor*. *Journal of Invertebrate Pathology*, 103, S96–S119.
- Ryabov, E.V., Wood, G.R., Fannon, J.M., Moore, J.D., Bull, J.C., Chandler, D., Mead, A., Burroughs, N. and Evans, D.J. (2014) A virulent strain of deformed wing virus (DWV) of honeybees (*Apis mellifera*) prevails after *Varroa destructor*-mediated, or in vitro transmission. *PLoS Pathogens*, 10, e1004230.
- Shimanuki, H., Calderone, N.W. and Knox, D.A. (1994) Parasitic mite syndrome—the symptoms. *American Bee Journal*, 134, 827–828.
- Sievers, F., Wilm, A., Dineen, D.G., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., Thompson, J.D. and Higgins, D. (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems Biology*, 7, e539.
- Singh, R., Levitt, A.L., Rajotte, E.G., Holmes, E.C. and Ostiguy, N. (2010) RNA viruses in Hymenopteran pollinators: evidence of inter-taxa virus transmission via pollen and potential impact on non-*Apis* Hymenopteran species. *PLoS ONE*, 5(12), e14357.
- Stevenson, M.A., Benard, H., Bolger, P. and Morris, R.S. (2005) Spatial epidemiology of the Asian honey bee mite (*Varroa destructor*) in the North Island of New Zealand. *Preventative Veterinary Medicine*, 71, 241–252.
- Tentcheva, D., Gauthier, L., Zappulla, N., Dainat, B., Cousserans, F., Colin, M.E. and Bergoin, M. (2004a) Prevalence and seasonal variations of six bee viruses in *Apis mellifera* L. and *Varroa destructor* mite populations in France. *Applied and Environmental Microbiology*, 70, 7185–7191.
- Tentcheva, D., Gauthier, L., Jouve, S., Canabady-Rochelle, L., Dainat, B., Cousserans F., Colin, M.E., Ball, B.V. and Bergoin, M. (2004b) Polymerase chain reaction detection of deformed wing virus (DWV) in *Apis mellifera* and *Varroa destructor*. *Apidologie*, 35, 431–440.
- Terio, V., Martella, V., Camero, M., Decaro, N., Testini, G., Bonerba, E., Tantillo, G. and Buonavoglia, C. (2008) Detection of a honeybee iflavirus with intermediate characteristics between kakugo virus and deformed wing virus. *New Microbiologica*, 31, 439–444.
- Topley, E., Davison, S., Leat, N. and Benjeddou, M., (2005) Detection of three honeybee viruses simultaneously by a single multiplex reverse transcriptase PCR. *African Journal of Biotechnology*, 4, 763–767.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. and Kumar, S. (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution*, 30, 2725–2729.
- vanEngelsdorp, D., Evans, J.D., Saegerman, C., Mullin, C. and Haubruge, E. (2009) Colony collapse disorder: a descriptive study. *PLoS ONE*, 4, e6481.
- vanEngelsdorp, D., Hayes, J.R.J., Underwood, R.M., Caron, D. and Pettis, J. (2011) A survey of managed honey bee colony losses in the USA, fall 2009 to winter 2010. *Journal of Apicultural Research*, 50, 1–10.
- vanEngelsdorp, D. and Meixner, M.D. (2010) A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *Journal of Invertebrate Pathology*, 103, S80–S95.
- Wang, H., Xie, J., Shreeve, T.G., Ma, J., Pallett, D.W. and King, L.A. (2013) Sequence recombination and conservation of *Varroa destructor* Virus-1 and deformed wing virus in field collected honey bees (*Apis mellifera*). *PLoS ONE*, 8, e74508.
- Williams, G.R., Rogers, R.E.L., Kalkstein, A.L., Taylor, B.A., Shutler, D. and Ostiguy, N. (2009) Deformed wing virus in

- western honey bees (*Apis mellifera*) from Atlantic Canada and the first description of an overtly-infected emerging queen. *Journal of Invertebrate Pathology*, 101, 77–79.
- Yang, X. and Cox-Foster, D.L. (2005) Impact of an ectoparasite on the immunity and pathology of an invertebrate: evidence for host immunosuppression and viral amplification. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 7470–7475.
- Yue, C. and Genersch, E. (2005) RT-PCR analysis of deformed wing virus in honeybees (*Apis mellifera*) and mites (*Varroa destructor*). *Journal of General Virology*, 86, 3419–3424.
- Zioni, N., Soroker, V. and Chejanovsky, N. (2011) Replication of *Varroa destructor* virus 1 (VDV-1) and a *Varroa destructor* virus 1-deformed wing virus recombinant (VDV-1-DWV) in the head of the honey bee. *Virology*, 417, 106–112.

Accepted September 14, 2015