

Biodegradation of diclofenac by activated sludge and membrane bioreactor-A review

D. Cherik, K. Louhab

Laboratory of Alimentary Technology, Faculty of Engineering Sciences,
 University of Boumerdes, Boumerdes 35000, Algeria

*Corresponding author: yandm96@yahoo.fr


ARTICLE INFO	ABSTRACT
Article History: Received : 01/01/2015 Accepted : 25/04/2015 Key Words: Biodegradation, Membrane bioreactor, Conventional activated sludge, Bacterial community	<i>Diclofenac (DCF) is a pharmaceutical residue of therapeutic class of non-steroidal anti-inflammatory which is often detected in the wastewater treatment plants (influent and effluent) and surface waters. This review focuses its elimination by biodegradation with activated sludge (CAS) or bioreactor membrane (MBR) in which microorganisms plays a key role in the elimination of diclofenac and in which a lot of factors can affect the efficiency of the removal as physicochemical properties of diclofenac, sludge retention time (SRT), temperature, pH, redox conditions and sludge characteristics. The objective of this study was to describe a review of the literature by recent publications on the biodegradation of diclofenac. We inspect the performance of biodegradation using biological process technology by activated sludge and membrane bioreactor in the elimination of diclofenac.</i>

I. Introduction

Diclofenac (DCF), a polycyclic non-steroidal anti-inflammatory drug [1- 3] is one of the most extensively studied pharmaceuticals, for his potential toxic effects on no-target organisms [4]. Diclofenac was detected in wastewater [5, 6] in a number of countries such as China [7, 8], Serbia [9], Spain [10, 11, and 12], Greece [13, 14], Portugal [15, 16], Brazil [17], USA [18] and South Africa [19]. According to published documents, the concentrations of this drug varied in a range from 0.01 to 8.5µg/l [20, 21] and the rate of his elimination in sewage treatment plants is low (up to 40%) [22, 23]. **Table 1** presents physicochemical properties and molecular structure of Diclofenac. It was considered very toxic anti-inflammatory due to the death of birds recorded shortly after scanning the infected farm in India and Pakistan [24, 25]; he showed potential influence of endocrine disruption by delaying and reducing the success of fish eggs and hatching low concentrations damage the digestive organs [25]. Sewage treatment plays an important part in removing contaminants from reclaimed water but it's known that conventional

wastewater treatment plants (WWTP) do not remove all the pollutants [26], especially the persistent polar pollutants due to their physicochemical properties such as diclofenac [27, 28, 29]. Most treatment methods of pharmaceuticals wastewater are biological processes [30, 14 and 31] in which microorganisms such as ammonia-oxidizing bacteria for the removal of ammonium [32], *Escherichia coli* [33] and nitrite-oxidizing bacteria [34] are applied for removal of organic contaminants [35, 25, and 36]. Most WWTPs used are activated sludge processes [37, 38] and membrane bioreactors [39, 40, and 24]. Biodegradation by these two processes can be influenced by various factors namely; the physicochemical properties of diclofenac [41], the biological operating conditions (sludge retention time, temperature, pH and redox conditions) [42-44, 45] and sludge characteristics [46, 47]. In this context, this study is a literature review of recent publications and studies to full and small scale on biodegradation of diclofenac to study the performance of activated sludge and membrane bioreactor in the elimination of diclofenac.

Table 1: Main characteristics of diclofenac

Structure _a	Molecular weight (g/mol) _a	Log K _{OW} _{a, b}	pK _a _a
	296.14	0.7-4.51	4.15

a : [48], b: [49]

II. Biodegradation of diclofenac

Biodegradation is the process most usually used to eliminate the pharmaceutical residues persistent as the diclofenac [50, 51 and 52, 53]. This process decomposes and mineralizes these pollutants by bacteria or mushrooms in simpler compounds [47, 54]. Diclofenac is a polar pharmaceutical compound mostly used as the sodium salt diclofenac-Na in human and veterinary medicine to reduce inflammation and pain [5, 17 and 55]. Many studies have evaluated the biodegradation of diclofenac by conventional activated sludge and membrane bioreactors.

2.1 Conventional activated sludge

Microbial degradation of diclofenac by activated sludge does not lead to its complete elimination but produces other metabolites which are also considered such as pollutants [56], these authors investigated the repartition of diclofenac in activated sludge, they detected 7 diclofenac transformation products among them 1-(2,6-dichlorophenyl)-1,3-dihydro-2H-indol-2-one and 2,3-dichloro-N-(phenyl)aniline. The fungus *Trametes versicolor* can be utilized in activated sludge for the elimination of diclofenac [57], in this study the removal rate was 64% which leads to a production of less toxic sludge, these authors concluded that fungal process can be an effective mean for biodegradation and diclofenac reduction in polluted waters. Activated sludge is the most used in the biological treatment and the rate of elimination of diclofenac by this process was insignificant and does not exceed 50 % [6, 43, 58, 59, 60, 61, 62 and 63]. Biodegradation of diclofenac can be estimated on measuring COD (Chemical Oxygen Demand) in polluted water and treated water [64], [60] have illuminated this point and studied the elimination of diclofenac by activated sludge suspended in 2 reactors; ASR1 and ASR2. The removal rate was 48±19% and the concentrations of DCO decreased from 976±39 mg/l and 707±14 mg/l to 47±48 mg/l and 54±55 mg/l in ASR1 and ASR2, respectively. The

hydrophobicity property defines the compound when the latter is insoluble in water [47], [65] studied the effect of the molecular features of diclofenac in activated sludge in lab-scale experimental; they confirm that the functional groups can influenced the biodegradability. Among the biological process conditions; the sludge retention time SRT which may have an effect on the biodegradation and also affects microbial activity [29], [51] evaluated the biodegradation of diclofenac in activated sludge of two wastewater treatment plants, Mamer and Boevange which differed in size, layout and sludge retention time (SRT). They observed that the biological degradation of diclofenac was significantly better in sludge from WWTP Mamer compared to sludge from WWTP Boevange and the removal rate decreases with increasing of SRT due to a lower active biomass presence. [66] reported the influence of SRT on diclofenac removal for more than seven months in denitrifying semi-continuous reactors operated at 10, 20, and 40 days by using two culture series (diclofenac free-control and diclofenac-acclimated); the results showed that the elimination rate for diclofenac-acclimated culture was lower than 15% a nitrate removal rate increased with increasing SRT but for diclofenac free-control this rate decreases with increasing SRT at 10, 20, 40 days with 48, 63, 79 mg NO₃⁻-N/L day and 122, 55, 54 mg NO₃⁻-N/L day, respectively. [67] investigated the biodegradation of diclofenac by nitrifying activated sludge and established the no removal of diclofenac with this process, these results are consistent with those of [68] who have studied the removal of diclofenac by nitrifying activated sludge in large scale (WWTP) and small scale (laboratory level) at a temperature of 12°C, the authors found that diclofenac is not biodegradable in large scale and the reaction rate constant (k') and biodegradation constant (k_{biol}) were 1.7 d⁻¹ and 0.6 l g_{ss}⁻¹d⁻¹, respectively in WWTP. In laboratory scale the biodegradation rate showed a difference between the reactors where with 10ug / l of diclofenac the biodegradation rate was 86% in the reactor 1, whereas in the other reactors was 90%. The authors explained this

difference by the amount of active nitrifying bacteria in the reactors which was influenced by temperature. [69] studied the removal of diclofenac in two lab-scale conventional activated sludge reactors under nitrifying (aerobic) and denitrifying (anoxic) conditions for more than 1.5 years, they found that the removal by the biological treatment with nitrifying or denitrifying bacteria was not significant (<25%). Diclofenac in anoxic conditions ($k_{\text{biol}} < 0.04 \text{ Lg}_{\text{ss}}^{-1}\text{d}^{-1}$) is persistent with respect to aerobic conditions ($k_{\text{biol}} = 1.2 \text{ Lg}_{\text{ss}}^{-1}\text{d}^{-1}$) [69]. [65] found a similar partition coefficient (k_d) under aerobic and anoxic conditions (0.319 and 0.303 L.gSS^{-1} , respectively).

Sludge characteristics may influence the sorption diclofenac. These characteristics vary depending on the operation of treatment plants as activated sludge [70]. [71] Studied the biodegradability of the diclofenac on two types of sludge; sterilized and activated. Results of batch adsorption experiments via sterilized sludge showed that the removal efficiency was 40.1% at 6 hours and 19.7% for activated sludge where the contributions of sludge adsorption and biodegradation were 14.9% at 6 hours and 4.8%, respectively, The authors suggested that this difference in removal efficiency by sterilized and activated sludge is due probably by the sterilizing effect of the activated sludge which can be caused changes in the characteristics of the sludge and improves the elimination of diclofenac and also suggested that the structure of diclofenac (the electronic withdrawing groups) namely its functional groups (amine, halogen, and carboxylic groups) causes the resistance of diclofenac in its biodegradation.

2.2 Membranes bioreactors

Recently, the membrane bioreactor process has become the most effective process for the removal of diclofenac compared to activated sludge [72], where the authors have demonstrated that the removal rate of diclofenac by membrane bioreactor process was 65%. The molecular features of diclofenac can affect the rate of his elimination by membrane bioreactor on the scale of laboratory [73]. In this study, the authors observed a very low rate of elimination (17 %) and they justified this behavior of diclofenac in the MBR system by the presence of chlorine group and also the electron withdrawing functional groups generates an electron deficiency [74] investigated the degradation of diclofenac by a white-rot fungus-augmented membrane bioreactor (MBR) with and without addition of a redox mediator (1-hydroxy benzotriazole, HBT); the results showed that the addition of a redox mediator (1-hydroxy

benzotriazole, HBT) improve the removal efficiency of diclofenac from 70% to 95%. On the other hand anaerobic MBR system has not shown an effective elimination rate for diclofenac (<10%) [75], these results are consistent with those of [76] these authors worked on a process of anaerobic / anoxic / aerobic membrane bioreactor in a large scale and observed removal rate below 20% for diclofenac. The full-scale MBR showed significantly better removal of hydrophilic compound ($\log D < 3$) [77]; they explained the greater removal by the existence of preand post-anoxic tanks and the combination of aerobic zones with different levels of DO (Dissolved Oxygen) relative to a pre-anoxic and one aerobic tank in the pilot MBR. In contrast in another study [28] where they used an anaerobic system by anaerobic fluidized membrane bioreactor (AFMBR) using granular activated carbon (GAC) as carrier medium, 78% was the removal rate of diclofenac and the elimination rate of COD was 95% at total HRT of 5 h. It can be concluded that the use of GAC helps eliminate diclofenac in the anaerobic system. In non-sterile conditions studied by [72] the removal rate of diclofenac was 55% in an MBR system with a hydraulic retention time of two days [78] have shown that increasing the biomass concentration in the MBR system due to the long retention time favors the elimination of diclofenac and can be influenced by microbial activity according to [79]. These authors found that the reduction of microbial activity in the MBR system induces a decrease in the rate of elimination of diclofenac which indicates that the latter is eliminated by biological degradation and not by the sorption process only. [80] investigated the elimination of diclofenac by MBR under different temperature (10-45°C) in laboratory-scale and demonstrated that the temperature has an effect on the microbial activity. The results showed that the maximum and minimum removal rate were at 10 ° C and 20°C, respectively.

As regards the influence of pH, [81] explain the relationship between physicochemical properties of diclofenac and his efficiency removal by a laboratory scale MBR at mixed liquor pH values of 5, 6, 7, 8, and 9. At pH 5 the rate of removal was highest; this result is due to the physicochemical properties of this ionisable compound under acidic conditions wherein the pK_a value is 4.15 [82] so at pH 5 diclofenac is neutral which makes it a hydrophobic compound. The increase of the made pH decreases $\log D$ and consequently the hydrophobicity of diclofenac what makes decrease also the efficiency elimination. [83] confirmed the results found by [81] with their study where they assessed the biotransformation of diclofenac by

laccase in an enzymatic membrane reactor, DCF biodegradation was better at acidic pH (3 and 4.5), and decreased with increasing pH.

3. Conclusion

The elimination of diclofenac was demonstrated to large and small scale with two process treatment of polluted water; biological process or biodegradation using conventional activated sludge (CAS) and membrane bioreactor (MBR) where the microorganisms are responsible for its degradation. Discussion of studies and publications used in this review demonstrated that biodegradation by membrane bioreactor is the most effective process for the removal of a resistant and persistent contaminant such as diclofenac with complete retention of suspended solids, thus reducing emissions to the dissolved fractions, but this removal may be influenced by various factors:

- Chemical structure where the presence of chlorine group and functional groups can influence the biodegradability of diclofenac.
- Sludge retention time (SRT) where the increase in this factor increasing the biomass concentration and therefore improves elimination of diclofenac.
- The temperature can affect the microbial activity in sludge and the maximum removal of diclofenac was seen at low temperatures.
- At acidic pH eliminating diclofenac was considered very favorable where the Hydrophobicity plays a very important role.
- Redox conditions: aerobic conditions in the elimination of diclofenac were better compared to the other conditions.

VI. References

1. Danina Krajišnik, Aleksandra Daković, Anđelija Malenović, Maja Milojević-Rakić, Vera Dondur, Željka Radulović, Jela Milić. Investigation of adsorption and release of diclofenac sodium by modified zeolites composites. *Applied Clay Science* 83–84 (2013) 322–326
2. ZHANG Nan, LIU HaiJin, LIU GuoGuan, WANG YingLing, LI Tong, WANG Gang. Role of cetyltrimethyl ammonium bromide, crystal violet and humic acid in the degradation of diclofenac under simulated sunlight irradiation. *SCIENCE CHINA Chemistry* December 2012 Vol.55 No.12: 2610–2616
3. Dorota Domaradzka, Urszula Guzik, Danuta Wojcieszynska. Biodegradation and biotransformation of polycyclic non-steroidal anti-inflammatory drugs. *Rev Environ Sci Biotechnol* (2015) 14:229–239
4. Abdul Jabbar Al-Rajab, Lyne Sabourin, David R. Lapen, Edward Topp. The non-steroidal anti-inflammatory drug diclofenac is readily biodegradable in agricultural soils. *Science of the Total Environment* 409 (2010) 78–82
5. Sung Kyu Maeng, Saroj K. Sharma, Karin Lekkerkerker-Teunissen, Gary L. Amy. Occurrence and fate of bulk organic matter and pharmaceutically active compounds in managed aquifer recharge: A review. *WATER RESEARCH* 45 (2011) 3015–3033
6. Jelena Radjenović, Mira Petrović, Damia Barceló. Fate and distribution of pharmaceuticals in wastewater and sewage sludge of the conventional activated sludge (CAS) and advanced membrane bioreactor (MBR) treatment. *WATER RESEARCH* 43 (2009) 831–841
7. Zhi-Hao Wen, Ling Chen, Xiang-ZhouMeng, Yan-PingDuan, Zeng-Sheng Zhang, EddyY.Zeng. Occurrence and human health risk of wastewater-derived pharmaceuticals in a drinking water source for Shanghai, East China.. *Science of the Total Environment* 490 (2014) 987–993
8. Guohua Dai, Bin Wang, Jun Huang, Rui Dong, Shubo Deng, Gang Yu. Occurrence and source apportionment of pharmaceuticals and personal care products in the Beiyun River of Beijing, China. *Chemosphere* 119 (2015) 1033–1039
9. Mira Petrović, Biljana Škrbić, Jelena Živančev, Laura Ferrando-Climent, Damia Barcelo. Determination of 81 pharmaceutical drugs by high performance liquid chromatography coupled to mass spectrometry with hybrid triple quadrupole–linear ion trap in different types of water in Serbia. *Science of the Total Environment* 468–469 (2014) 415–428
10. Dolores Camacho-Muñoz, Julia Martín, Juan Luis Santos, Irene Aparicio, Esteban Alonso. Concentration evolution of pharmaceutically active compounds in raw urban and industrial wastewater. *Chemosphere* 111 (2014) 70–79
11. M.D. Camacho-Munoz, J.L. Santos, I. Aparicio, E. Alonso. Presence of pharmaceutically active compounds in Donana Park (Spain) main watersheds. *Journal of Hazardous Materials* 177 (2010) 1159–1162
12. Emma Gracia-Lor, Juan V. Sancho, Roque Serrano, Félix Hernández. Occurrence and removal of pharmaceuticals in wastewater treatment plants at the Spanish Mediterranean area of Valencia. *Chemosphere* 87 (2012) 453–462
13. Christina I. Kosma, Dimitra A. Lambropoulou, Triantafyllos A. Albanis. Investigation of PPCPs in wastewater treatment plants in Greece: Occurrence, removal and environmental risk assessment. *Science of the Total Environment* 466–467 (2014) 421–438
14. Christina I. Kosma, Dimitra A. Lambropoulou, Triantafyllos A. Albanis. Occurrence and removal of PPCPs in municipal and hospital wastewaters in Greece. *Journal of Hazardous Materials* 179 (2010) 804–817
15. Aleksandar Lolić, Paula Paíga, Lúcia H.M.L.M. Santos, Sandra Ramos, Manuela Correia, Cristina Delerue-Matos. Assessment of non-steroidal anti-inflammatory and analgesic pharmaceuticals in seawaters of North of Portugal: Occurrence and environmental risk. *Science of the Total Environment* 508 (2015) 240–250
16. André M.P.T. Pereira, Liliana J.G. Silva, Leonor M. Meisel, Celeste M. Lino. Environmental impact of pharmaceuticals from Portuguese wastewaters: geographical and seasonal occurrence, removal and risk assessment. *Environmental Research* 136 (2015) 108–119
17. Aldo Pacheco Ferreira. Trace Analysis of Pharmaceutical Residues in Wastewater Treatment Plants in Rio de Janeiro, Brazil. *Archive of SID. Journal of Chemical Health Risks* (2014) 4 (1), 1–12
18. Tiago S. Oliveira, Mark Murphy, Nicholas Mendola, Virginia Wong, Doreen Carlson, Linda Waring. Characterization of Pharmaceuticals and Personal Care products in hospital effluent and waste water influent/effluent by direct-injection LC-MS-MS. *Science of the Total Environment* 518–519 (2015) 459–478
19. Foluso O. Agunbiade & Brenda Moodley. Pharmaceuticals as emerging organic contaminants in

- Umgeni River water system, KwaZulu-Natal, South Africa. *Environ Monit Assess* (2014) 186:7273–7291
20. Lúcia H.M.L.M. Santos, A.N. Araújo, Adriano Fachini, A. Pena, C. Delerue-Matos, M.C.B.S.M. Montenegro. Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. *Journal of Hazardous Materials* 175 (2010) 45–95
21. M. D. Vedenyapina, D. A. Borisova, A. P. Simakova, L. P. Proshina, and A. A. Vedenyapin. Adsorption of Diclofenac Sodium from Aqueous Solutions on Expanded Graphite. ISSN 0361-5219, *Solid Fuel Chemistry*, 2013, Vol. 47, No. 1, pp. 59–63. © Allerton Press, Inc., 2013.
22. J. Hartmann, P. Bartels, U. Mau, M. Witter, W.v. Tümpling, J. Hofmann, E. Nietzschmann. Degradation of the drug diclofenac in water by sonolysis in presence of catalysts. *Chemosphere* 70 (2008) 453–461
23. Jia-Qian Jiang, Z. Zhou, V.K. Sharma. Occurrence, transportation, monitoring and treatment of emerging micro-pollutants in waste water — A review from global views. *Microchemical Journal* 110 (2013) 292–300
24. J.L. Zhou, Z.L. Zhang, E. Banks, D.Grover, J.Q. Jiang. Pharmaceutical residues in wastewater treatment works effluents and their impact on receiving river water. *Journal of Hazardous Materials* 166 (2009) 655–661
25. Asu Ziyilan, Nilsun H. Ince. The occurrence and fate of anti-inflammatory and analgesic pharmaceuticals in sewage and fresh water: Treatability by conventional and non-conventional processes. *Journal of Hazardous Materials* 187 (2011) 24–36
26. Chenxi Wu, Xiaolong Huang, Jason D. Witter, Alison L. Sponberg, Kexiong Wang, Ding Wang, Jiantong Liu. Occurrence of pharmaceuticals and personal care products and associated environmental risks in the central and lower Yangtze river, China. *Ecotoxicology and Environmental Safety* 106 (2014) 19–26
27. P. Verlicchi, M. Al Aukidy, E. Zambello. Occurrence of pharmaceutical compounds in urban wastewater: Removal, mass load and environmental risk after a secondary treatment—A review. *Science of the Total Environment* 429 (2012) 123–155
28. Kasturi Dutta, Ming-Yi Lee, Webber Wei-Po Lai, Chien Hsien Lee, Angela Yu-Chen Lin, Cheng-Fang Lin, Jih-Gaw Lin. Removal of pharmaceuticals and organic matter from municipal wastewater using two-stage anaerobic fluidized membrane bioreactor. *Bioresource Technology* 165 (2014) 42–49
29. Yasemin Kaya, Gamze Ersan, Ilda Vergili, Z. Beril Gönder, Gulsum Yilmaz, Nadir Dizge, Coskun Aydinler. The treatment of pharmaceutical wastewater using in a submerged membrane bioreactor under different sludge retention times. *Journal of Membrane Science* 442 (2013) 7282
30. Ioanna A. Vasiliadou, Raul Molina, Fernando Martínez, Juan A. Melero. Biological removal of pharmaceutical and personal care products by a mixed microbial culture: Sorption, desorption and biodegradation. *Biochemical Engineering Journal* 81 (2013) 108–119
31. Aleksandra Jelic, Meritxell Gros, Antoni Ginebreda, Raquel Cespedes-Sánchez, Francesc Ventura, Mira Petrovic, Damià Barceló. Occurrence, partition and removal of pharmaceuticals in sewage water and sludge during wastewater treatment. *WATER RESEARCH* 45 (2011) 1165–1176
32. Shuyi Wang, Claudia K. Gunsch. Effects of selected pharmaceutically active compounds on the ammonia oxidizing bacterium *Nitrosomonas europaea*. *Chemosphere* 82 (2011) 565–572
33. Laure Pasquini, Christophe Merlin, Lucille Hassenboehler, Jean-François Munoz, Marie-Noëlle Pons, Tatiana Görner. Impact of certain household micropollutants on bacterial behavior. Toxicity tests/study of extracellular polymeric substances in sludge. *Science of the Total Environment* 463–464 (2013) 355–365
34. Barbara Kraigher, Ines Mandic-Mulec. Nitrification activity and community structure of nitrite-oxidizing bacteria in the bioreactors operated with addition of pharmaceuticals. *Journal of Hazardous Materials* 188 (2011) 78–84
35. José Rivera-Utrilla, Manuel Sánchez-Polo, María Ángeles Ferro-García, Gonzalo Prados-Joya, Raúl Ocampo-Pérez. Pharmaceuticals as emerging contaminants and their removal from water. A review. *Chemosphere* 93 (2013) 1268–1287
36. Ling-Wei Meng, Xiang-kun Li, Ke Wang, Kai-Li Ma, Jie Zhang. Influence of the amoxicillin concentration on organics removal and microbial community structure in an anaerobic EGSB reactor treating with antibiotic wastewater. *Chemical Engineering Journal* 274 (2015) 94–101
37. Miao Jiang, Weiben Yang, Ziwei Zhang, Zhen Yang, Yuping Wang. Adsorption of three pharmaceuticals on two magnetic ion-exchange resins. *JOURNAL OF ENVIRONMENTAL SCIENCES XX* (2015) XXX–XXX
38. Ai Jia, Yi Wan, Yang Xiao, Jianying Hu. Occurrence and fate of quinolone and fluoroquinolone antibiotics in a municipal sewage treatment plant. *WATER RESEARCH* 46 (2012) 387–394
39. Kok Kwang Ng, Xueqing Shi, Yinuo Yao, How Yong Ng. Bio-entrapped membrane reactor and salt marsh sediment membrane bioreactor for the treatment of pharmaceutical wastewater: Treatment performance and microbial communities. *Bioresource Technology* 171 (2014) 265–273
40. Noor Sabrina Ahmad Mutamim, Zainura Zainon Noor, Mohd Ariffin Abu Hassan, Gustaf Olsson, Adhi Yuniarto. Membrane bioreactor: Applications and limitations in treating high strength industrial wastewater. *Chemical Engineering Journal* 225 (2013) 109–119
41. Luong N. Nguyen, Faisal I. Hai, Shufan Yang, Jinguo Kang, Frederic D.L. Leusch, Felicity Roddick, William E. Price, Long D. Nghiem. Removal of pharmaceuticals, steroid hormones, phytoestrogens, UV-filters, industrial chemicals and pesticides by *Trametes versicolor*: Role of biosorption and biodegradation. *International Biodeterioration & Biodegradation* 88 (2014) 169–175
42. [42] Meritxell Gros, Mira Petrović, Antoni Ginebreda, Damià Barceló. Removal of pharmaceuticals during wastewater treatment and environmental risk assessment using hazard indexes. *Environment International* 36 (2010) 15–26
43. Sonia Suárez, Rubén Reif, Juan M. Lema, Francisco Omil. Mass balance of pharmaceutical and personal care products in a pilot-scale single-sludge system: Influence of T, SRT and recirculation ratio. *Chemosphere* 89 (2012) 164–171
44. Eleni N. Evgenidou, Ioannis K. Konstantinou, Dimitra A. Lambropoulou. Occurrence and removal of transformation products of PPCPs and illicit drugs in wastewaters: A review. *Science of the Total Environment* 505 (2015) 905–926
45. Shengnan Xu, Donglei Wu, Zhiqiang Hu. Impact of hydraulic retention time on organic and nutrient

- removal in a membrane coupled sequencing batch reactor. *WATER RESEARCH* 55 (2014) 12–20
46. Gianpaolo Sabia, Marco Ferraris, Alessandro Spagni. Effect of solid retention time on sludge filterability and biomass activity: Long-term experiment on a pilot-scale membrane bioreactor treating municipal wastewater. *Chemical Engineering Journal* 221 (2013) 176–184
 47. Magdalena Cirja, Pavel Ivashechkin, Andreas Schäffer, Philippe F. X. Corvini. Factors affecting the removal of organic micropollutants from wastewater in conventional treatment plants (CTP) and membrane bioreactors (MBR). *Rev Environ Sci Biotechnol* (2008) 7:61–78
 48. Kangmin Chon, Jaewon Cho, Ho Kyong Shon. A pilot-scale hybrid municipal wastewater reclamation system using combined coagulation and disk filtration, ultrafiltration, and reverse osmosis: Removal of nutrients and micropollutants, and characterization of membrane foulants. *Bioresource Technology* 141 (2013) 109–116
 49. Evangelos C. Symaris, Ioannis A. Fotidis, Athanasios S. Stasinakis, Irini Angelidaki. Effects of triclosan, diclofenac, and nonylphenol on mesophilic and thermophilic methanogenic activity and on the methanogenic communities. *Journal of Hazardous Materials* 291 (2015) 45–51
 50. A. I. Rodarte-Morales, G. Feijoo, M. T. Moreira, J. M. Lema. Biotransformation of three pharmaceutical active compounds by the fungus *Phanerochaete chrysosporium* in a fed batch stirred reactor under air and oxygen supply. *Biodegradation* (2012) 23:145–156
 51. Marius Majewsky, Tom Gallé, Viviane Yargeau, Klaus Fischer. Active heterotrophic biomass and sludge retention time (SRT) as determining factors for biodegradation kinetics of pharmaceuticals in activated sludge. *Bioresource Technology* 102 (2011) 7415–7421
 52. Ngoc Han Tran, Taro Urase, Huu Hao Ngo, Jiangyong Hu, Say Leong Ong. Insight into metabolic and cometabolic activities of autotrophic and heterotrophic microorganisms in the biodegradation of emerging trace organic contaminants. *Bioresource Technology* 146 (2013) 721–731
 53. Hyun-Jeoung Lee, Eunhwa Lee, Sung-Hwa Yoon, Hee-Ra Chang, Kyun Kim, Jung-Hwan Kwon. Enzymatic and microbial transformation assays for the evaluation of the environmental fate of diclofenac and its metabolites. *Chemosphere* 87 (2012) 969–974
 54. Aida García-Rodríguez, Víctor Matamoros, Clàudia Fontàs, Victòria Salvadó. The ability of biologically based wastewater treatment systems to remove emerging organic contaminants—a review. *Environ Sci Pollut Res* (2014) 21:11708–11728
 55. Anna Barra Caracciolo, Edward Topp, Paola Grenni. Pharmaceuticals in the environment: Biodegradation and effects on natural microbial communities. A review. *Journal of Pharmaceutical and Biomedical Analysis* 106 (2015) 25–36
 56. Tina Kosjek, Ester Heath, Sandra Pérez, Mira Petrović, Damià Barceló. Metabolism studies of diclofenac and clofibric acid in activated sludge bioreactors using liquid chromatography with quadrupole – time-of-flight mass spectrometry. *Journal of Hydrology* 372 (2009) 109–117
 57. Carlos E. Rodríguez-Rodríguez, Aleksandra Jelic, Marta Llorca, Marinella Farré, Mira Petrovic, Damià Barceló, Teresa Vicent, Glòria Caminal. Solid-phase treatment with the fungus *Trametes versicolor* substantially reduces pharmaceutical concentrations and toxicity from sewage sludge. *Bioresource Technology* 102 (2011) 5602–5608
 58. Bruce Petrie, Ewan J. McAdam, Mark D. Scrimshaw, John N. Lester, Elise Cartmell. Fate of drugs during wastewater treatment. *Trends in Analytical chemistry* 49 (2013) 145–159
 59. Jia-Qian Jiang, Z. Zhou, V.K. Sharma. Occurrence, transportation, monitoring and treatment of emerging micro-pollutants in waste water — A review from global views. *Microchemical Journal* 110 (2013) 292–300
 60. P. Falås, P. Longrée, J. la Cour Jansen, H. Siegrist, J. Hollender, A. Joss. Micropollutant removal by attached and suspended growth in a hybrid biofilm-activated sludge process. *WATER RESEARCH* 47 (2013) 4498–4506
 61. Luong N. Nguyen, Faisal I. Hai, William E. Price, Frederic D.L. Leusch, Felicity Roddick, Hao H. Ngo, Wenshan Guo, Saleh F. Magram, Long D. Nghiem. The effects of mediator and granular activated carbon addition on degradation of trace organic contaminants by an enzymatic membrane reactor. *Bioresource Technology* 167 (2014) 169–177
 62. Mojca Zupanc, Tina Kosjek, Martin Petkovšek, Matevž Dular, Boris Kompare, Brane Širok, Željko Blažeka, Ester Heath. Removal of pharmaceuticals from wastewater by biological processes, hydrodynamic cavitation and UV treatment. *Ultrasonics Sonochemistry* 20 (2013) 1104–1112
 63. R. Salgado, R. Marques, J. P. Noronha, G. Carvalho, A. Oehmen, M. A. M. Reis. Assessing the removal of pharmaceuticals and personal care products in a full-scale activated sludge plant. *Environ Sci Pollut Res* (2012) 19:1818–1827
 64. G. Buttiglieri, T.P.Knepper. Removal of Emerging Contaminants in Wastewater Treatment: Conventional Activated Sludge Treatment. *Hdb Env Chem Vol. 5, Part S/2* (2008): 1–35
 65. M. Pomiès, J. M. Choubert, C. Wisniewski, C. Miège, H. Budzinski, M. Coquery. Lab-scale experimental strategy for determining micropollutant partition coefficient and biodegradation constants in activated sludge. *Environ Sci Pollut Res* (2015) 22:4383–4395
 66. Gamze Ozdemir, Egemen Aydin, Emel Topuz, Cigdem Yangin-Gomec, Didem Okutman Tas. Acute and chronic responses of denitrifying culture to diclofenac. *Bioresource Technology* 176 (2015) 112–120
 67. P. Falås, A. Baillon-Dhumez, H.R. Andersen, A. Ledin, J. la Cour Jansen. Suspended biofilm carrier and activated sludge removal of acidic pharmaceuticals. *WATER RESEARCH* 46 (2012) 1167–1175
 68. Antonina Kruglova, Pia Ahlgren, Nasti Korhonen, Pirjo Rantanen, Anna Mikola, Riku Vahala. Biodegradation of ibuprofen, diclofenac and carbamazepine in nitrifying activated sludge under 12 °C temperature conditions. *Science of the Total Environment* 499 (2014) 394–401
 69. Sonia Suarez, Juan M. Lema, Francisco Omil. Removal of Pharmaceutical and Personal Care Products (PPCPs) under nitrifying and denitrifying conditions. *WATER RESEARCH* 44 (2010) 3214–3224
 70. Li Wang, Jian Zhang, Ran Zhao, Cong Li, Ye Li, Chenglu Zhang. Adsorption of basic dyes on activated carbon prepared from *Polygonum orientale* Linn: Equilibrium, kinetic and thermodynamic studies. *Desalination* 254 (2010) 68–74
 71. Huiju Fan, Jing Li, Liqiu Zhang, Li Feng. Contribution of sludge adsorption and biodegradation to the removal of five pharmaceuticals in a submerged membrane bioreactor. *Biochemical Engineering Journal* 88 (2014) 101–107
 72. Shufan Yang, Faisal I. Hai, Long D. Nghiem, Luong N. Nguyen, Felicity Roddick, William E. Price. Removal of bisphenol A and diclofenac by a novel fungal membrane bioreactor operated under non-sterile conditions. *International Biodeterioration & Biodegradation* 85 (2013) 483–490

73. Nichanan Tadkaew, Faisal I. Hai, James A. McDonald, Stuart J. Khan, Long D. Nghiem. Removal of trace organics by MBR treatment: The role of molecular properties. *WATER RESEARCH* 45 (2011) 2439-2451
74. Luong N. Nguyen, Faisal I. Hai, Shufan Yang, Jinguo Kang, Frederic D.L. Leusch, Felicity Roddick, William E. Price, Long D. Nghiem. Removal of trace organic contaminants by an MBR comprising a mixed culture of bacteria and white-rot fungi. *Bioresource Technology* 148 (2013) 234-241
75. Victor M. Monsalvo, James A. McDonald, Stuart J. Khan, Pierre Le-Clech. Removal of trace organics by anaerobic membrane bioreactors. *WATER RESEARCH* 49 (2014) 103-112
76. Wenchao Xue, Chunying Wu, Kang Xiao, Xia Huang, Haidong Zhou, Hiroshi Tsuno, Hiroaki Tanaka. Elimination and fate of selected micro-organic pollutants in a full-scale anaerobic/anoxic/aerobic process combined with membrane bioreactor for municipal wastewater reclamation. *WATER RESEARCH* 44 (2010) 5999-6010
77. Hop V. Phan, Faisal I. Hai, James A. McDonald, Stuart J. Khan, Ren Zhang, William E. Price, Andreas Broeckmann, Long D. Nghiem. Nutrient and trace organic contaminant removal from wastewater of a resort town: Comparison between a pilot and a full scale membrane bioreactor. *International Biodeterioration & Biodegradation* xxx (2015) 1-9
78. E. Fernandez-Fontaina, F. Omil, J.M. Lema, M. Carballa. Influence of nitrifying conditions on the biodegradation and sorption of emerging micropollutants. *WATER RESEARCH* 46 (2012) 5434-5444
79. Sung Kyu Maeng, Byeong Gyu Choi, Kyu Tae Lee, Kyung Guen Song. Influences of solid retention time, nitrification and microbial activity on the attenuation of pharmaceuticals and estrogens in membrane bioreactors. *WATER RESEARCH* 47 (2013) 3151-3162
80. Faisal I. Hai, Karin Tessmer, Luong N. Nguyen, Jinguo Kang, William E. Price, Long D. Nghiem. Removal of micropollutants by membrane bioreactor under temperature variation. *Journal of Membrane Science* 383 (2011) 144-151
81. Nichanan Tadkaew, Muttucumaru Sivakumar, Stuart J. Khan, James A. McDonald, Long Duc Nghiem. Effect of mixed liquor pH on the removal of trace organic contaminants in a membrane bioreactor. *Bioresource Technology* 101 (2010) 1494-1500
82. Kangmin Chon, Jaeweon Cho, Ho Kyong Shon. A pilot-scale hybrid municipal wastewater reclamation system using combined coagulation and disk filtration, ultrafiltration, and reverse osmosis: Removal of nutrients and micropollutants, and characterization of membrane foulants. *Bioresource Technology* 141 (2013) 109-116
83. Luong N. Nguyen, Faisal I. Hai, William E. Price, Frederic D.L. Leusch, Felicity Roddick, Ewan J. McAdam, Saleh F. Magram, Long D. Nghiem. Continuous biotransformation of bisphenol A and diclofenac by laccase in an enzymatic membrane reactor. *International Biodeterioration & Biodegradation* 95 (2014) 25-32

Please cite this Article as:

Cherik, D., Louhab, K., Biodegradation of diclofenac by activated sludge and membrane bioreactor- A review, , ***Algerian J. Env. Sc. Technology*, 1:1 (2015) 11-17**